RETINA ASSOCIATES OF KENTUCKY ONE WEEK AFTER SURGERY INSTRUCTIONS

Name	Date

You had retina surgery last week. The first week is usually the most difficult time, but there are still ways you can help make the surgery a success.

Eye Medication

In most cases, you will need to continue with Neo/Poly/Dex ointment. You will taper off of this.

i	NOC 24208 795-35 BAUSCH + LOMB
	Neomycin and Polymyxin B
	Sulfates and Dexamethasone
	Ophthalmic Ointment USP
	(Stetle)
п	*
п	
	(mirror)
п	3.5,00
	Tidiilla (i)

Neo/Poly/Dex- Use ¼ inch strip at bedtime for 1 wee	ek
then stop.	
Additional Drops:	

What can I do (activity limits) after the first week?

You can begin to resume normal activities, but for another week, don't do anything that causes a lot of strain. Still avoid getting water in the eye. You no longer need the shield at night. Ask your doctor if you have questions about a specific activity you wish to do. Usually, two weeks after surgery you can go back to your regular activities.

What position should I be in?

If you had to be in a special position the first week, your doctor will tell you how much longer he wants you to stay that way.

Your Position Should Be (circled):

FACE DOWN	RIGHT SIDE DOWN	LEFT SIDE DOWN
UPRIGHT	ANY POSITION	NOT ON YOUR BACK
OTHER	FOR HOW LONG:	

What should I expect?

Any swelling, pain and redness should continue to improve. If they don't, call us. How soon you will see is different for everyone, but it can take a month or longer. Wait until your one month visit with us before you go back to your regular eye doctor for new glasses.

What should I do if there is a problem or concern?

If you have a question or concern, call us anytime, at **1-800-627-2020** and ask for the doctor on call. Thank you for trusting us with your vision.

William Wood, MD Rick Isernhagen, MD Thomas Stone, MD John Kitchens, MD Todd Purkiss, MD Belinda Shirkey, MD Sheila Garcia Santana, MD Blake Isernhagen, MD